


MOLECULAR MACHINERY: A Tour of the Protein Data Bank

Living cells are filled with complex molecular machinery, a million times smaller than familiar machines like computers or automobiles. Cells use these tiny molecular machines to perform all of the jobs needed for life. Some are molecular scissors that cut food into cell-sized pieces. Some build new molecules when cells grow or when damaged tissues are repaired. Some are molecular bones and muscles that support cells and help them move and crawl. Some fight off attackers, defending against infection.

Researchers around the world are studying these molecules and determining their precise atomic structures. These structures are available on the Internet through the Protein Data Bank (<http://www.pdb.org>), the central storehouse of biomolecular structures. A few of the thousands of structures held in the Protein Data Bank are shown here. In these pictures, the molecules are all drawn at a magnification of 3,000,000 times, and each atom is shown as a small sphere. Many of these structures are composed of several subunits, which are indicated by different colors.

An enormous range of sizes is shown here: the water molecule at the left has only three atoms and the rhinovirus shown below has hundreds of thousands.

By David S. Goodsell, The Scripps Research Institute, La Jolla, California, USA
Graphic design by Gail W. Bamber, San Diego Supercomputer Center



PDB PROTEIN DATA BANK

<http://www.pdb.org/> • info@rcsb.org
RESEARCH COLLABORATORY FOR
STRUCTURAL BIOINFORMATICS

RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY
NEW YORK UNIVERSITY
NEW YORK UNIVERSITY
NATIONAL INSTITUTE OF STANDARDS AND TECHNOLOGY

All proteins in the original and current versions of this figure are available in PDB format. • **PDB ID: 1a00:** Insulin. • **PDB ID: 1a01:** Glucagon. • **PDB ID: 1a02:** Human Growth Hormone. • **PDB ID: 1a03:** Interferon. • **PDB ID: 1a04:** Trypsin. • **PDB ID: 1a05:** Amylase. • **PDB ID: 1a06:** Phospholipase. • **PDB ID: 1a07:** Lysozyme. • **PDB ID: 1a08:** Ribonuclease. • **PDB ID: 1a09:** Pepsin. • **PDB ID: 1a10:** Antibody. • **PDB ID: 1a11:** Cycloxygenase. • **PDB ID: 1a12:** Cytchrome c. • **PDB ID: 1a13:** Cytochrome c Oxidase. • **PDB ID: 1a14:** Cytchrome bcl Complex. • **PDB ID: 1a15:** Aquaporin. • **PDB ID: 1a16:** Potassium Channel. • **PDB ID: 1a17:** Mechanosensitive Channel. • **PDB ID: 1a18:** Gramicidin. • **PDB ID: 1a19:** Porin. • **PDB ID: 1a20:** Ferritin. • **PDB ID: 1a21:** Serum Albumin. • **PDB ID: 1a22:** Rhodopsin. • **PDB ID: 1a23:** Photosynthetic Reaction Center. • **PDB ID: 1a24:** Light-Harvesting Complex. • **PDB ID: 1a25:** Cytidine Triphosphate (ATP).

• **PDB ID: 1a26:** Topoisomerase. • **PDB ID: 1a27:** RNA Polymerase. • **PDB ID: 1a28:** Valyl-tRNA Synthetase. • **PDB ID: 1a29:** Threonyl-tRNA Synthetase. • **PDB ID: 1a30:** Isoleucyl-tRNA Synthetase. • **PDB ID: 1a31:** Phenylalanyl-tRNA Synthetase. • **PDB ID: 1a32:** Aspartyl-tRNA Synthetase. • **PDB ID: 1a33:** Glutamyl-tRNA Synthetase. • **PDB ID: 1a34:** Catalytic Gene Activator Protein. • **PDB ID: 1a35:** TATA-binding Protein/Transcription Factor II B. • **PDB ID: 1a36:** DNA Polymerase. • **PDB ID: 1a37:** 70S Ribosome. • **PDB ID: 1a38:** 40S Transfer RNA. • **PDB ID: 1a39:** Elongation Factor Tu. • **PDB ID: 1a40:** Elongation Factor Tu and Ts. • **PDB ID: 1a41:** Elongation Factor G. • **PDB ID: 1a42:** Proline cis/trans Isomerase. • **PDB ID: 1a43:** Prefoldin. • **PDB ID: 1a44:** Chaperonin GroEL/ES. • **PDB ID: 1a45:** Microtubule. • **PDB ID: 1a46:** Myosin. • **PDB ID: 1a47:** Actin. • **PDB ID: 1a48:** Collagen.

2 MEMBRANES

Cells are surrounded by a membrane made of lipids, like the phospholipid and cholesterol molecules shown at the top. Membranes keep the cellular machinery inside and unwanted material out. Many proteins are embedded in this membrane, performing a variety of essential tasks. ATP synthase is one of the rotary generators that produces ATP (adenosine triphosphate), the small molecule used for powering cells. The two large complexes below it change a battery that powers ATP synthase, and the tiny protein cytochrome c shuttles electrons between them. Rhodopsin is found in membranes in the retina. The small retinol molecule inside it changes shape when illuminated, causing the surrounding protein to send a signal to the brain. Cyclooxygenase builds one of the molecules used to signal pain—this cyclooxygenase molecule here, however, is blocked by two molecules of aspirin, shown inside in white. At the bottom are three molecules involved in photosynthesis, which capture energy from light and use it to power the synthesis of sugar in plant cells.

3 TRANSPORT AND STORAGE

Of course, a perfectly sealed membrane would be of little use to cells, because nutrients could not get in and wastes could not get out. The box shows a membrane looking face-on. Five proteins that form channels through the membrane are shown. To the right of the box are several soluble proteins involved in transport and storage of molecules. Hemoglobin and myoglobin carry oxygen. Ferritin forms a hollow shell that stores iron ions. Serum albumin carries many different molecules in the blood.

4 CHEMICAL FACTORIES

Cells build a bewildering variety of enzymes—proteins that perform chemical reactions. At the top are the ten enzymes that perform glycolysis, the breakdown of sugar to form ATP. Below that are several enzymes that perform different housekeeping reactions. Dihydrofolate reductase activates a key cofactor molecule and alcohol dehydrogenase breaks down alcohol. Ribulose bisphosphate carboxylase/oxygenase is the most common enzyme on the Earth, and performs a key step in the capture of carbon dioxide by plants to form sugar. The three synthases and the transferase make different building blocks for creating new molecules. Nitrogenase performs an essential role in the ecosystem by converting nitrogen gas into a form that living cells can use.

5 DNA

Genetic information is stored in the DNA double helix, seen running from top to bottom here. Many proteins are used to copy, read, and store this information. RNA polymerase copies the information into a strand of RNA that will be used to direct the construction of new proteins. It is assisted by topoisomerase, which releases tension when the helix is wound and unwound, and guides to appropriate starting points by the two protein complexes below it. DNA polymerase replicates DNA strands—here, the polymerase is filling a gap in the double helix. Some proteins, like the lac repressor, grab DNA and bend it sharply, or even wrap it all the way around themselves, like the two nucleosomes at the bottom.

6 BUILDING NEW PROTEINS

New proteins are built by ribosomes—complex molecular factories that read the genetic code and use it to direct construction. Many molecular machines are needed to assist the process. Twenty different aminoacyl-tRNA synthases (six are shown here) load the building blocks onto tRNA, ready to be added to a growing protein chain. Several protein factors, shown below the ribosome, guide each tRNA into the proper spot. The three chaperone proteins shown at the bottom help each new protein fold into its proper shape.

7 BEAMS AND GIRDERS

Cells are braced and supported by a complex infrastructure. This cytoskeleton is formed of sturdy filaments like actin and microtubules, composed of many subunits stacked like bricks. Myosin is a molecular motor that climbs along actin filaments, allowing the cell to move. Collagen, broken into two pieces here, is actually found outside of cells, where it forms connective tissue between cells.

